



TITLE:

On Morin Singularities(C^∞ - maps and their Singularities)

AUTHOR(S):

Ando, Yoshifumi

CITATION:

Ando, Yoshifumi. On Morin Singularities(C^∞ - maps and their Singularities). 数理解析研究所講究録 1985, 550: 126-130

ISSUE DATE:

1985-02

URL:

<http://hdl.handle.net/2433/98883>

RIGHT:

On Morin Singularities

by

Yoshifumi Ando

(安藤 良文)

Introduction

In this paper we study the problem of finding a smooth map between smooth manifolds with nice Morin singularities in a given homotopy class. A geometric interpretation of Morin singularities of a smooth map $f:N \rightarrow P$ is as follows. Let $S^i(f)$ denote the set of all points x of N such that the kernel rank of df_x is i . For a certain map f , $S^i(f)$ becomes a submanifold of N and we may define $S^{i,j}(f)$ as the set $S^j(f|S^i(f))$ for $f|S^i(f):S^i(f) \rightarrow P$ similarly. Let $n = \dim N$, $p = \dim P$ and $i = \max(1, n-p+1)$. Let I_r be the r -sequence $(i, 1, \dots, 1)$. Then we may continue to define $S^{I_r}(f)$ as $S^1(f|S^{I_{r-1}})$ inductively. A point of $S^{i,0}(f)$ or $S^{I_r}(f)$ is called a Morin singularity of symbol $(i,0)$ or I_r respectively. However this approach does not make it clear for what part of smooth maps f , $S^{I_r}(f)$ can be defined. For this we review the following important observation due to Boardman[2].

There exist a submanifold $\Sigma^{i,0}(N,P)$ and a series of submanifolds; $\Sigma^{I_1}(N,P) \supset \Sigma^{I_2}(N,P) \supset \dots \supset \Sigma^{I_r}(N,P) \supset \dots$ in the infinite jet space $J^\infty(N,P)$. The codimension of $\Sigma^{i,0}(N,P)$ is $i(p-n+i)$ and that of $\Sigma^{I_r}(N,P)$ is $n-p+r$ for $n \geq p$ and

$r(p-n+1)$ for $n < p$. He has shown that if a jet map $j^\infty f: N \rightarrow J^\infty(N, P)$ of f is transverse to all submanifolds $\Sigma^{i,0}(N, P)$ and $\Sigma^{I_r}(N, P)$, then $S^{i,0}(f)$ and $S^{I_r}(f)$ coincide with $(j^\infty f)^{-1}(\Sigma^{i,0}(N, P))$ and $(j^\infty f)^{-1}(\Sigma^{I_r}(N, P))$ respectively. Therefore for generic maps f we may consider $S^{i,0}(f)$ and $S^{I_r}(f)$.

For any integer $r \geq 1$ we define a subset $\Omega_r(N, P)$ of $J^\infty(N, P)$ as the set of all jets z such that either z is of maximal rank or a point of $\Sigma^{i,0}(N, P)$ or $\Sigma^{I_2}(N, P) \setminus \Sigma^{I_{r+1}}(N, P)$. Then $\Omega_r(N, P)$ becomes an open subbundle of the fibre bundle $J^\infty(N, P)$ over N . The first result of this paper is the following

Theorem 1. Let $p \geq 2$. Then for any section s of N into $\Omega_r(N, P)$, there exists a smooth map $g: N \rightarrow P$ such that $j^\infty g$ becomes a section of N into $\Omega_r(N, P)$ homotopic to s in $\Omega_r(N, P)$.

Next we will study the problem of eliminating the Morin singularities $S^{I_r}(f)$ with $\text{codim } S^{I_r}(f) = n$ from f admitting only Morin singularities. Theorem 1 reduces it to a problem of finding a section of N into $\Omega_{r-1}(N, P)$ homotopic to $j^\infty f$. We will show that if $j^\infty f$ is transverse to $\Sigma^{I_r}(N, P)$ for a connected and closed manifold N , then the number of points of $S^{I_r}(f)$ modulo 2 is the unique obstruction of finding the above section. We should note that this number is just the Thom polynomial of the topological closure $\overline{\Sigma^{I_r}(N, P)}$ for f (see the definition of [9]).

Theorem 2. Let $r \geq 2$, $p \geq 2$ and $\text{codim } \Sigma^{I_r}(N, P) = n$. Let N and P be orientable manifolds. Then

- (1) A smooth map f with $j^\infty f(N) \subset \Omega_r(N, P)$ is homotopic to

a smooth map g such that $j^\infty g(N) \subset \Omega_{r-1}(N, P)$ and $j^\infty f$ and $j^\infty g$ are homotopic as sections of N into $\Omega_r(N, P)$ if and only if the Thom polynomial of $\overline{\Sigma^I_r}(N, P)$ for f vanishes.

(2) In particular f is homotopic to such a smooth map g in the following cases;

- i) $n > p$ and $r \equiv 1 \pmod{4}$
- ii) $n > p$, $r \equiv 2, 3$ or $4 \pmod{4}$ and $n-p \equiv 1 \pmod{2}$ and
- iii) $n \leq p$ and $n+p+r+\frac{1}{2}r(r+1) \equiv 0 \pmod{2}$.

It will be shown by the Morse inequalities that the similar statement of Theorem 1 for $p = 1$ is not true. If N is an open manifold, then Theorem 1 is a direct consequence of Gromov[7, Theorem 4.1.1] and if $n < p$, it is also a special case of [4, Theorem B]. So the rest cases will be treated in this paper. The case $r = 2$ of Theorem 1 should be compared with [6, Theorem 1.3] which will play an important role in a proof of Theorem 1 (Sections 2 and 3).

The case $n \geq p$ and $p = 2$ of Theorem 2 has been proved by Levine[11, Theorems 1 and 2] for $n > 2$ and by Eliasberg[5, Corollary of Theorem 4.9] for $n = 2$.

References

- [1] Y. Ando, On the elimination of certain Thom-Boardman singularities of order two, *J. Math. Soc. Japan*, 34(1982), 241-268.
- [2] J. M. Boardman, Singularities of differentiable maps, *IHES Publ. Math.*, 33(1967), 21-57.
- [3] J. Damon, Thom polynomials for contact class singularities, thesis, Harvard Univ., 1972.
- [4] A. A. du Plessis, Maps without certain singularities, *Comment. Math. Helv.*, 50(1975), 363-382.
- [5] J. M. Eliasberg, On singularities of folding types, *Math. USSR Izv.*, 4(1970), 1119-1134.
- [6] J. M. Eliasberg, Surgery of singularities of smooth mappings, *Math. USSR Izv.*, 6(1972), 1302-1326.
- [7] M. L. Gromov, Stable mappings of foliations into manifolds, *Math. USSR Izv.*, 3(1969), 671-694.
- [8] M. W. Hirsch, Immersions of manifolds, *Trans. Amer. Math. Soc.*, 93(1959), 242-276.
- [9] A. Haefliger and A. Kosinski, Un théorème de Thom sur les singularités des applications différentiables, *Séminaire H. Cartan E.N.S.*, 1956/57, Exposé N°8.
- [10] A. Lascoux, Calcul de certains polynômes de Thom, *C.R. Acad. Sc. Paris*, 278(1974), 889-891.
- [11] H. I. Levin, Elimination of cusps, *Topology* 3(1965), 263-296.
- [12] S. Łojasiewicz, Triangulation of semianalytic sets *Ann. Scuola. Norm. Sup. Pisa. Sci. Fis. Mat.*, 18(1959), 87-136.

- [13] B. Morin, Formes canoniques des singularités d'une application différentiable, C. R. Acad. Sc. Paris, 260 (1965), 5662-5665, 6503-6506.
- [14] F. Ronga, Le calcul des classes duales singularités de Boardman d'ordre deux, Comment. Math. Helv., 47 (1972), 15-35.
- [15] F. Ronga, Le calcul de la classe de cohomologie entière duale à Σ^k , Springer Lecture Notes, 192 (1971), 313-315.
- [16] F. Sergeraert, Expression explicite de certains polynômes de Thom, C. R. Acad. Sc. Paris, 276 (1973), 1661-1663.
- [17] N. Steenrod, The Topology of Fibre Bundles, Princeton Univ. Press, Princeton, 1951.
- [18] R. Thom, Les singularités des applications différentiables, Ann. Inst. Fourier, 6 (1955-56), 43-87.